

**A Study on Anemia and Its Association with Hematological Parameters in Different Age Groups**Pramod Shinde¹, P Jogi Naidu², Shamili Moningi³¹Associate Professor, Department of Pathology, Great Eastern Medical School& Hospital, Srikakulam, India.²Assistant Professor, Department of Pathology, Great Eastern Medical School& Hospital, Srikakulam, India.³Assistant Professor, Department of Pathology, Great Eastern Medical School& Hospital, Srikakulam, India.**Correspondence Author:** Pramod Shinde, Associate Professor, Department of Pathology, Great Eastern Medical School& Hospital, Srikakulam, India.**Conflicts of Interest:** Nil**Abstract**

Objective: To examine the occurrence of anemia and assess the hematological parameters of subjects concerned and study the type of anemia in various age groups.

Methods: This study was conducted in Great Eastern Medical School& Hospital, Srikakulam in March 2016. Total 450 cases were studied. The cases were inside patient admitted in different wards in Great Eastern Medical School& Hospital, Srikakulam. The sample for test was collected in EDTA tube. The slides were prepared and stained with field stain. The sample was run in hematology cell counter Sysmex 3 part for hematology indices and other parameters. Microscopic examination of slides was done for peripheral smear examination and complete blood count estimation.

Results: The overall prevalence of anemia was 42.44%. Mild degree of Anemia was present in 46.15%, which was the most common followed by moderate degree which was present in 43.58% and severe anemia was present in 10.25%, which was the least common of degree of anemia. Anemia was present in 51.05% of men and in 48.95% of women. Microcytic hypochromic anemia is the most common type of anemia in overall age groups 46.66%. Normocytic normochromic anemia without anisocytosis was the most common type of

anemia 42.05% in adult patient. Mild degree of anemia is more common in males 51.32% and moderate and severe anemia was more in females 49.23% and 50.07% respectively.

Conclusion: The incidence of anemia increases with age also allied with race, chronic diseases, nutritional deficiencies and other circumstances such as infection. In diverse age group the incidence of various types of anemia is different, which is because of diverse etiology and severity in this age groups. As a result, a diagnosis of anemia warrants enough clinical attention, to find out the cause, type, severity and this forms the basis for treatment of anemia.

Keywords: Anemia, Hematology, indices, Iron deficiency, Chronic diseases.

Introduction

Anemia is defined as a decrease of the total circulating red cell mass below normal limits. Anemia reduces the oxygen-carrying capacity of the blood, foremost to tissue hypoxia. In put into practice, the measurement of red cell mass is not easy, and anemia is typically diagnosed based on a decrease in the hematocrit (the ratio of packed red cells to total blood volume) and the hemoglobin concentration of the blood to levels that are below the normal range.^[1]

There are many classifications of anemia and the clinical manifestations of the anemia are nonspecific and were detailed earlier. The signs and symptoms are gastrointestinal or gynecologic disease, malnutrition, pregnancy, and malabsorption.^[1] The diagnosis of iron deficiency anemia ultimately rests on laboratory studies.

Anemia of Chronic Disease ^[1]

According to the World Health Organization (WHO), there are two billion people with anemia in the world and half of the anemia is due to iron deficiency ^[2]. Anemia is a late indicator of iron deficiency, so it is estimated that the prevalence of iron deficiency is 2.5 times that of anemia ^[2]. With 40 per cent prevalence of anemia in the world on an average for the general population, the prevalence in the developing countries tends to be three to four times higher than in the developed countries.

However anemia should not be accepted as an inevitable consequence of ageing^[5]. Studies indicate that the prevalence of anemia increases with advancing age and under age 75 years, anemia is more frequent in females, but over age 75 years it is more frequent in males^[6]. Multiple pathophysiologic abnormalities in a single elderly patient with anemia are well known. Micronutrient deficiencies as reason of anemia have been frequently documented in the aged people.^[7]

WHO reported that, anemia is because of different pathophysiological mechanisms. The most prevalent types of anemia are due to nutritional deficiencies (malnutrition and iron, vitamin B12 and folic acid deficiencies) and chronic diseases (such as cancer, kidney disease and congestive heart failure)^[8,10]. In order to characterize the type of anemia and formulate a differential diagnosis, the work-up should include physical exams and laboratory tests, such as evaluations

of hematocrit, hemoglobin and red blood cell indices. The red blood cell indices should include the cell count, MCV, mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC) and red cell distribution width (RDW)^[11]. In fact, the hemoglobin concentration is the factor that is most frequently used as an indicator of the pathophysiological consequences of anemia. However, this variable is not very specific or sensitive. Hemoglobin levels can be altered in different pathologic conditions, such as infectious and inflammatory processes, hemorrhage, protein-caloric malnutrition, associated to medications and smoking. The MCV guides the diagnosis of anemia and helps in its classification. However, the MCV value, that is, the mean size of the red cells (macrocytic, microcytic and normocytic), should be used together with the RDW, thus directing the interpretation of the variation in the size of red blood cells.

Aims: To assess the hematological parameters of subjects involved, analyze the prevalence, type and severity of spectrum of anemia in various age groups.

Methodology

The study was carried out in Great Eastern Medical School & Hospital, Srikakulam. The patients of all age groups admitted in various wards of hospital were included in study. Outdoor patient were excluded from study. The selection was performed randomly. The blood samples were collected in EDTA tubes and were immediately sent to the laboratory for hematological testing. The hematological testing was performed in the Hematology division of the Clinical Pathology Laboratory of Great Eastern Medical School & Hospital, Srikakulam. The equipment used included a Sysmax 3part analyzer. The evaluated parameters included the hemoglobin concentration and red blood cell indices- Mean Cell Volume (MCV), Mean cell hemoglobin

(MCH), Mean cell hemoglobin concentration (MCHC), hematocrit (PCV), Red blood cell count, total leucocyte count, differential count and platelet count.

Heamoglobin concentrations (g/dL) For the diagnosis of anaemia and assessment of severity according to the World Health Organization.			
Age	Mild	Moderate	Severe
6-59 months	10-10.9	7-9.9	<7
5-11 years	11-11.4	8-10.9	<8
12-14 years	11-11.9	8-10.9	<8
Female >14 years	11-11.9	8-10.9	<8
Male >14 years	11-12.9	8-10.9	<8

The reference range of Mean Cell Volume (MCV) was taken as 80-100fl, mean cell hemoglobin (MCH) was taken as 27-32pg, for mean cell hemoglobin concentration 32-36g/dl. Microcytic anemia was taken as MCV value less than 80fl and MCH less than 27. Macrocytic was taken when MCV is greater than 100fl. Normocytic Normochromic was taken when all hematological indices are within range.

Observation & Results

According to a UNICEF report 2 billion people suffer from anemia worldwide and most of them have IDA, especially in underdeveloped/developing countries.

According to WHO, almost 20 % of [21] all women of the childbearing age in United States were suffering from iron deficiency anemia as compared to 2 % of adult males. According [22] to WHO, in the developing countries, about 50-60 % of young children and pregnant females and 20-30 % non-pregnant females were affected by iron deficiency anemia. In countries where little meat is in the diet, [22] iron deficiency anemia is 6-8 times more prevalent than in North America. Thus it is one of the global problems, mainly affecting the [23] developing countries. Pregnant and lactating females, growing children and elderly people with some underlying disease

causing blood loss are at more risk as compared to other groups of population.

Anaemia is the most prevalent nutritional deficiency disorder in the world. It affects all age groups. Globally, anaemia affects 1.62 billion people, which corresponds to 24.8% of the population. The highest prevalence of anemia exists in the developing world where its causes are multi-factorial. National Family Health Survey statistics reveal that every second Indian woman is anaemic and one in every five maternal deaths is directly due to anaemia.

This study having total 450 cases. The cases were inside patient admitted in various wards in Great Eastern Medical School& Hospital, Srikakulam.

Table 1 – Prevalence of Anemia

	Total number of Subjects	Anemic Subjects
Number of Subjects	450	195
Percentage	100%	43.33%

Out of 450 studied patient anemia was found 195 patient of different age group. Prevalance of anemia in indoor patient is 43.33%.

Table 2 Gender distribution of subjects

	Total Numbers	Percentage
Males	99	50.07
Females	96	49.23

In the study total number of males were 99 and female were 96. There was no statistical difference in both gender. Mild anemia was common in males 50.07% , while moderate and severe anemia was more common in females.

Table 3 – Grading of Anemia

Severity of Anemia	Numbers of subjects	Percentage
Mild	90	46.15%
Moderate	85	43.58%
Severe	20	10.25%
Total	195	100%

In our study, Mild anemia was most common followed by moderate and severe anemia.

Table 4 – Gender wise Distribution of Grading of Anemia

	Mild	Moderate	Severe
Male	47 (52.21%)	42 (49.01%)	10 (48.33%)
Female	43 (47.79%)	42 (50.99%)	10 (51.67%)
	91 (100%)	84 (100%)	20 (100%)

Table 5 – Age wise Distribution of Grading of Anemia

Age in years	Mild	Moderate	Severe	Total	Percentage
>40	43	33	8	84	43.07
31 – 40	17	16	04	37	18.97
21 – 30	12	18	04	34	17.43
11 – 20	14	08	02	24	12.30
<10	04	08	03	15	7.69

Anemia is most common in adult population, 43.10%, second peak was seen in the age group 31-40(19.08%). In our study anemia was least common in age group <10 years.

Table-6 Peripheral smear examination in Mild anemia

Peripheral smear examination	Total subjects	Percentage
Normocytic normochromic	66	73.33%
Microcytic hypochromic	22	24.44%
Normocytic normochromic, few Macrocytes	2	2.22%

Table-7 Peripheral smear examination in Moderate anemia

Peripheral smear examination	Total subjects	Percentage
Normocytic normochromic	16	18.82
Microcytic hypochromic	55	67.07
Macrocytic normochromic	14	17.07

Table-8 Peripheral smear examination in Severe anemia

Peripheral smear examination	Total subjects	Percentage
Normocytic normochromic	1	5.2
Microcytic hypochromic	13	68.42
Macrocytic normochromic	03	15.78
Dimorphic	02	10.52

Table-9 Prevalance of anemia in population

Peripheral smear examination	Total subjects	Percentage
Normocytic normochromic	82	42.05
Microcytic hypochromic	91	46.66
Macrocytic normochromic	19	9.74
Dimorphic	3	1.53

Microcytic Hypochromic Group:

Out of 450 patients 91 had microcytic hypochromic anemia (46.66%). This is the most common type of anemia in the study. Majority of the subjects (55) had moderate degree of anemia and 22 had mild type followed by severe type of anemia, which is seen in 14 subjects.

Normocytic Normochromic Group

Out of 450 patients 82 had normocytic normochromic anemia accounting for 42.05% of anemic patients. 73.33% of Normocytic group had mild anemia. Majority in this group were adult males.

Macrocytic Normochromic Group

Out of 450 patients 19 had macrocytic normochromic anemia accounting for 9.74% of anemic patients. of

Normocytic group had mild anemia. 13 patient in this group (15.78%) had moderate anemia, 3 had severe anemia, and 2 had mild anemia. Dimorphic group: Out of 450 patients had 2 dimorphic anemia. All in this group had severe anemia (100%).

Discussion

In this study using routine clinical data from a large number of patients, anaemia was present in the majority of adults and young children.

The high proportion of microcytic anaemia and the fact that gender differences were only seen after the menarche period in women indicate that iron deficiency was the main cause of anaemia. In a study of children aged 12–23 months in two rural districts in India, 72% of children with anaemia had low ferritin levels .[8] Other Indian studies have also shown high prevalence of iron deficiency anaemia among young women [9, 10]. The high prevalence of iron deficiency anaemia among women in childbearing age has important public health implications. It is estimated that anaemia accounts for 12.8% of maternal mortality in Asia [11]. Iron requirements are greater in pregnancy, and iron deficiency is associated with maternal death, preterm delivery, and low birth-weight [12, 13]. In India, only 28% of women consume meat, fish, or eggs on a weekly basis [4] and the iron bioavailability of the vegetarian diet is poor. [10, 14] Effective public health programmes aimed at reducing iron deficiency among young women could have a major impact in reducing maternal and infant mortality. [15]

Literature has revealed that ageing does have an effect on blood production with reduced ratio of bone marrow to fat cells and reduced marrow response when stimulated with erythropoietin. [7] However, the decline of hemoglobin and resulting increase in anemia with age should not be presumed to be a result of "normal aging" and blanket treatment with hematinics should be avoided. Literature

has revealed that ageing does have an effect on blood production with reduced ratio of bone marrow to fat cells and reduced marrow response when stimulated with erythropoietin. MCV increases slightly with increasing age but usually not enough to produce significant macrocytosis. Although there is a paradoxical feedback in renal production of erythropoietin, since the levels of this hormone actually increase over time, it has also been reported that the erythroid marrow may become less sensitive to erythropoietin stimulation, a key factor contributing along with possible nutritional deficits and comorbidities to the development of anaemia in the elderly. Even distinguishing anemia of chronic inflammation from anemia of chronic kidney disease is somewhat challenging considering the fact that increased inflammation is seen in older adults even without chronic kidney disease and there are coexisting morbidities in this age group. [16] Anemia algorithms used for evaluation of younger adults are based on the mean corpuscular volume. Such algorithms may be less helpful in the elderly because the classic changes in erythrocyte size do not often accompany anemia in this age group. In most elderly patients with anemia, red cell indices disclose normocytic, normochromic anemia. Clinicians therefore might begin the evaluation of anemia as they would in younger adults, but, if they do not find one of the classic causes of microcytosis or macrocytosis, the search for a cause might need to be enlarged. [17]

Anemia is a critical clinical problem in the elderly population with a significant public health impact. Data on the prevalence of anemia are varied and depend on the location and the population. The present study demonstrated a prevalence of anemia of 43.48%. This high percentage of prevalence is similar to the S.Patel et al study.[17] The prevalence of anemia in the highest in

adults (>40 years) which is 42.9%.The similarities in prevalence may be associated with a greater concern with the aging population and the greater emphasis on their health. The increase in the frequency of anemia in older individuals is sometimes regarded as inherent to the aging process. Nevertheless, further investigations of the cause of anemia and the completion of treatment may help to improve clinical conditions in the elderly population. It is important to remember that anemia is multifactorial, and its occurrence may be due to the presence of cancer, inflammatory diseases, kidney disease (due to diabetes and hypertension), and the use of several drugs commonly required in the elderly population.[18] Our findings demonstrate a higher prevalence of anemia in women than in men considering all the age groups. These results are in similar to those of a study that evaluated 284 elderly participants in the Family Health Program of Pernambuco, Brazil. The study found that the average prevalence of anemia in women was 12.6% versus 11.8% in men. In contrast, a study performed by Olivares et al.[19] enrolled 274 elderly outpatients in Chile and found a low prevalence of overall anemia, but a higher prevalence among men (5.4%) compared to women (4.4%). This increase in the prevalence of anemia in men was also described in the NHANES III study, with men accounting for 11.6% of the cases of anemia versus 10.2% for women. [5] In an attempt to explain the different prevalence rates of anemia for men and women, some authors have argued that estrogens act as inhibitors of erythropoiesis and make women more vulnerable to the development of anemia. However, while postmenopausal estrogen levels decrease, there is an increase in red cell mass to levels that are similar to those in males, which makes it unreasonable to use different criteria for anemia in each gender.

[20, 21] In our study 99(50.07%) were males and 96(49.23%) were females, which was similar to the Kaur et al[24] in which 37% were males and 33% were females, and in contrast to the Chul won choi et al study in which 11.4% were males and 2.1% were females.[25] In our study Microcytic hypochromic anemia 46.66% is the predominant type of anemia as seen in Gerardo et al studies [26] and S Patel et al [27] study in which microcytic hypochromic anemia was seen in 72%.This finding was in contrast to the Kaur et al [24] in which normocytic normochromic anemia is the predominant type 56%.

In our study normocytic normochromic anemia was found in 42.05%, majority of which had mild type of anemia 73.52%. The predominant age group in this category was elderly.

In our study the majority of the subjects were adults (43.07%), followed by the patient in the third decade (18.97%), this was similar to the Kaur et al study[24] in which 55% were found in the age group 60-69 years, and in contrast to the S. Patel et al study[27] in which peak age group was 21-30 years,46%.

Conclusion

The occurrence of anemia in patients admitted in our hospital was higher than population based studies. Hematological parameters guiding the type of anemia diverge in assorted age groups involved which reflect the varying etiologies behind this. In younger age group the main type is microcytic hypochromic which is due to iron deficiency. In adults the type most prevalent is normocytic normochromic which may be because of chronic diseases, inflammation, blood loss, malignancies or aging process. However, anemia is not a state that should only be linked with the aging process. Anemia is not a condition, it is a symptom of a variety of pathologies which deserves adequate therapeutic awareness.

References

- [1]. Kumar, Abbas, Fausto, Aster, Robins and Cotran, Pathologic Basis of Disease, Eight edition, Chapter 14.
- [2]. WHO, UNICEF, and UNU, Iron Deficiency Anaemia: Assessment, Prevention and Control, A Guide for Programme Managers, WHO, UNICEF, UNU, Geneva, Switzerland, 2001, http://www.who.int/nutrition/publications/micronutrients/anaemia_iron_deficiency/WHO_NHD_01.3/en/index.html.
- [3]. R. D. Baker, F. R. Greer, and Committee on Nutrition American Academy of Pediatrics, –Diagnosis and prevention of iron deficiency and iron-deficiency anemia in infants and young children (0–3 years of age), *Journal of Pediatrics*, vol. 126, pp. 1040–1050, 2010.
- [4]. World Health Organisation, Iron Deficiency Anaemia Assessment, Prevention, and Control, A Guide for Programme Managers, WHO, 2001.
- [5]. World Health Organization. Definition of an older or elderly person. Retrieved August 29, 2010. <http://www.who.int/healthinfo/survey/ageingdefnolder/en/index.html>.
- [6]. Ferrucci L, Semba RD, Guralnik JM, Ershler WB, Bandinelli S, Patel KV et al. Proinflammatory state, hepcidin and anemia in older persons. *Blood*. 2010;115:3810-26.
- [7]. Russell RM, Rasmussen H, Fada RD. The Impact of Nutritional Needs of Older Adults on Recommended Food Intakes. *Nutrition in Clinical Care* 1999;2:164–76.
- [8]. S.-R. Pasricha, J. Black, S. Muthayya et al., –Determinants of anemia among young children in rural India, *Journal of Pediatrics*, vol. 126, no.1, pp. e140–e149, 2010.
- [9]. K. C. Menon, S. A. Skeaff, C. D. Thomson et al., –Concurrent micronutrient deficiencies are prevalent in nonpregnant rural and tribal women from central India, *Journal of Nutrition*, vol. 27, no. 4, pp. 496–502, 2011.
- [10]. P. Thankachan, S. Muthayya, T. Walczyk, A. V. Kurpad, and R. F. Hurrell, –An analysis of the etiology of anemia and iron deficiency in young women of low socioeconomic status in Bangalore, India, *Food and Nutrition Bulletin*, vol. 28, no. 3, pp. 328–336, 2007.
- [11]. K. S. Khan, D. Wojdyla, L. Say, A.M. Gulmezoglu, and P. F. van Look, –WHO analysis of causes of maternal death: a systematic review, *The Lancet*, vol. 367, no. 9516, pp. 1066–1074, 2006.
- [12]. K. Kalaivani, –Prevalence & consequences of anaemia in pregnancy, *Indian Journal of Medical Research*, vol. 130, no. 5, pp. 627–633, 2009.
- [13]. L. H. Allen, –Anemia and iron deficiency: effects on pregnancy outcome, *The American Journal of Clinical Nutrition*, vol. 71, no. 5, pp. 1280s–1284s, 2000.
- [14]. K. Shridhar, P. K. Dhillon, L. Bowen et al., –Nutritional profile of Indian vegetarian diets—the Indian Migration Study (IMS), *Nutrition Journal*, vol. 13, article 55, 2014.
- [15]. Z. A. Bhutta, J. K. Das, R. Bahl et al., –Can available interventions end preventable deaths in mothers, newborn babies, and stillbirths, and at what cost? *The Lancet*, vol. 384, no. 9940, pp. 347–370, 2014.
- [16]. Beutler E, Waalen J. The definition of anemia: what is the lower limit of normal of the blood hemoglobin concentration? *Blood* 2006;107:1747.
- [17]. Chaves PH, Xue QL, Guralnik JM. What constitutes normal hemoglobin concentration in community-dwelling disabled older women? *J Am Geriatr Soc* 2004;52:1811.
- [18]. Cliquet MG. Como diagnosticar e tratar anemia no idoso. *Rev Bras Med*. 2010;67(4):89-9619.
- [19]. Olivares M, Hertramp E, Capurro MT, Wegner D. Prevalence of anemia in elderly subjects living at home: role of micronutrient deficiency and inflammation. *Eur J Clin Nutr*. 2000; 54(11):834-9.

- [20]. Failace R. Hemograma: manual de interpretação. Porto Alegre: Artmed; 1995. p.197.
- [21]. Sahadevan S, Choo PW, Jayaratnam FJ. Anaemia in the hospitalized elderly. Singapore Med J. 1995;36(4):375-8.
- [22]. World Health Organization. Iron deficiency anaemia: assessment, prevention and control. A guide for programme managers. WHO: Geneva; 2001.
- [23]. Gualandro SF, Hojaij NH, Jacob Filho W. Deficiência de ferro no idoso. Rev Bras Hematol Hemoter. 2010;32(supl. 2):57-61.
- [24]. Kaur H, Piplani S, Madan M. Prevalence of anemia and micronutrient deficiency in elderly. International Journal of Medical and Dental Science. 2014;3(1):296-302.
- [25]. Choi CW, Lee J, Park KH, Yoon SY, Choi IK, Oh SC. Prevalence and Characteristics of Anemia in the Elderly: Cross –Sectional Study of Three Urban Korean Population Samples. Am J Hematol 2004;77(1):26-30.
- [26]. Gerardo Alvarez Uria, Praveen K. Naik, Manoranjan Midde, Pradeep S. Yalla, Raghavakalyan Pakam, Prevalance and severity of anaemia stratified by age and gender in rural India, Hindawi Publishing Corporation, Anemia 2014.
- [27]. S. Patel, M. Shah, J. Patel, N. Kumar, Iron Deficiency anemia in moderate to severely anemic patients, Gujarat Medical Journal, August-2009, vol.64, No.2;15-18.