

International Journal of Medical Science and Innovative Research (IJMSIR) IJMSIR : A Medical Publication Hub

Available Online at: www.ijmsir.com

Volume - 3, Issue -1, January - 2018, Page No. : 237 - 244

**Biological Activities of Talaromyces Purpurogenus T013** 

Giang Vo, Hien Tran, Anh Nguyen, Yen Nguyen, Tu Nguyen\*

School of Biotechnology, HoChiMinh City International University, Vietnam National University – Ho Chi Minh City,

Vietnam, Quarter 6, Linh Trung ward, Thu Duc district, Hochiminh city, Vietnam

Correspondence Author: Tu Nguyen

Type of publication: Original Research Paper

## **Conflicts of Interest:** Nil

### Abstract

In the study, Talaromyces purpurogenus T013, a filamentous fungi belonging to Talaromyces which was identified had been isolated from the high value Taxus baccata subsp. wallichiana. In order to find out a new source of antimicrobial and anticancer agents, T. purpurogenus cultured in YM was against five different kinds of pathogens including Salmonella typhi (16.77±0.21 mm), Pseudomonas aeruginosa (16.17±0.21 mm), Serratia marcescens (15.50±0.21 mm), Listeria monocytogenes (16.37±0.15 and Vibrio mm) parahaemolyticus (15.43±0.21 mm) by using the agar well diffusion method. There was cytotoxic ability on Hela cancer cell line at the concentration (100 mg/mL) by using sulforhodamin B assay. The chloroform fraction showed antimicrobial activities that suggested the purified agent with antimicrobial in this fraction will be obtained so far.

**Keywords:** *Talaromyes, cytotoxicity, antimicrobial activities*, chloroform fraction.

## 1. Introduction

A famous plant produces taxol in Vietnam is *Taxus* baccata subsp. wallichiana. Endophytic fungi isolated from *Taxus baccata subsp. wallichiana* have the ability to produce biological compounds [1].

*Talaromyces purpurogenus (T. purpurogenus)* produces rubratoxins [2] [3] [4]. Rubratoxin B causes mutation, hepatotoxicity to several animals [5] [6] [7] [8] [9]. The first report of rubratoxicosis to human was reported by Richer et al. (1997) [10]. *T. purpurogenus* could produce gluconic acid. However, it could produce potential antitumor agent [11] [12].

At present, antimicrobial resistance caused by *Pseudomonas aeruginosa*, *Salmonella typhi*, *Listeria monocytogenes*, *Vibrio parahaemolyticus* and others [13] [14] [15] [16].

Therefore, this study tried to detect the biological activities of *T. purpurogenus* isolated from *Taxus baccata subsp. wallichiana* in order to find out a new source of antimicrobial and anticancer agents production that serve for resistance in the future. *T. purpurogenus* is expected to have the antimicrobial activities on different pathogens such as *Salmonella typhi*, *Pseudomonas aeruginosa*, *Serratia marcescens*, *Listeria monocytogenes* and *Vibrio parahaemolyticus*. The purification step using polar solvent and non-polar solvent is able to collect the pure substance that has the antimicrobial ability with above pathogens and cytotoxicity on Hela cells [17] [18].

### 2. Materials and methods

## 2.1. Fungi collection and screening

The trunk, leaves were separated and cultured in PDB (Potato Dextro broth) for enrichment. After 4-5 days, the culture was spread onto PDA containing *Bacillus subtilis* as indicator. The colonies inhibited *Bacillus subtilis* were

transferred on YM medium containing 1% glucose, 0.3% yeast extract, 0.3% malt extract, 0.5% peptone and 2% agar which is used for maintenance and sporulation of fungi.

## 2.2. Antimicrobial test

Pathogens used for antimicrobial test were *Salmonella typhi* ATCC 19430, *Pseudomonas aeruginosa* ATCC 27853 (P3), *Serratia marcescens* ATCC13880 (P6), *Listeria monocytogenes* [17] and *Vibrio parahaemolyticus* (VSA). All these pathogens were cultured in Luria-Bertani (LB) broth and gram-stained to check the purity. Before the test, the pathogen samples were measured the absorbance to determine the optical density (OD). The test was performed on LB agar plate [19]. The supernatant extracted from *T. purpurogenus* culture will be used as the sample of the test.

The antimicrobial test was performed using agar well diffusion method [20]. M1 medium was used as the negative control for the test while tetracycline was used as the positive control. Applying aseptic techniques, using a small sterile metal stick to punch into the LB agar plate to form wells. Amount (10<sup>6</sup> CFU/mL) of pathogen cultures (Salmonella typhi, Pseudomonas aeruginosa, Serratia Listeria monocytogenes Vibrio marcescens, and parahaemolyticus) was spread onto the surface of the plate, using sterile swabs. Finally, the supernatant was pipetted into the wells. All the plates were incubated at room temperature. After ten to twelve hours, the inhibition zones formed on the wells were measured and analyzed.

### **2.3.** Cytotoxicity test

Sulforhodamine B and different cancer cell lines (HepG2, MCF-7, Hela and NCI-H460) were used for the test. The sample prepared for this test was the dry powder extracted from *T* .*purpurogenus* culture mixed with EMEM/10%

FBS medium which was used for culture of cancer cell lines.

Positive control of the test was the mixture of cells and camptothecin at the concentration 0.07  $\mu$ g/ml. The mixture of cells and the solvent that dissoved the *T. purpurogenus* dry powder (EMEM/10% FBS culture medium) was used as negative control. The test for the sample at each concentration was designed with four wells. Two wells contained the cancer cell line, the culture (EMEM/ 10% FBS) and the tested substance (*T. purpurogenus* dry powder), the other two contained only the culture and the sample (blank).

The cytotoxicity test was performed by sulforhodamine B assay [21]. It is a simple colorimetric method for determining sensitivity and cytotoxicity of a substance. Sulforhodamine B (SRB), a negatively charged dye which can bind electrostatically with the positively charged parts of proteins will be the main reagent for the assay. The amount of binding dye will reflect the total cellular protein.

Firstly, the cancer cell line preserved in liquid nitrogen will be thawed and cultured until the fourth generation (P4). Next, the cells will be cultured in culture flasks to achieve the coverage of about 70-80%. The cells now is covered into wells of 96-well plate at a density of  $10^4$  cells per well (In the case of HepG2, MCF-7 and Hela) or 7.5 x  $10^3$  cells per well (In the case of NCI-H460 cancer cell line). After that, incubate the cell at  $37^{\circ}$ C in 5% CO<sub>2</sub> for 24 hours. After incubation, the wells are supplied with the medium of the tested substance with the double concentration. Continue to incubate the cells at  $37^{\circ}$ C in 5% CO<sub>2</sub> for 48 hours. The cells in the wells now are fixed with trichloroacetic acid (TCA).

Secondly, cells were dyed with SRB: 0.2% SRB was loaded into each well and incubated at room temperature (25°C) for 5 to 20 minutes. After that, SRB was discarded

and the cells were washed gently with 1% acetic acid for 5 times. Let the cells dry at room temperature  $(25^{\circ}C)$  from 12 hours to 24 hours. The results were obtained by loading 200 µL Tris-base into each well, then shaking for 10 to 15 minutes until the SRB was dissolved completely and measuring the optical density value at wavelength 492 nm and 620 nm.

The cytotoxic percentage was calculated by the formula:

$$\%I = \left(1 - \frac{OD_{sample}}{OD_{control}}\right) \times 100\%$$

Notes:

OD<sub>Cell</sub>: OD value of well containing cells

OD<sub>Blank</sub>: OD value of blank well

OD<sub>sample</sub>: OD value of sample obtained from formula (1) and (2)

OD<sub>control</sub>: OD value of control obtained from formula (1) and (2)

### 2.4. Preliminary purification

Supernatant extracted from T. purpurogenus culture after 240h of culturing will be used for purification. To purify the supernatant, chloroform and petroleum ether were used as non-polar solvents and methanol as polar solvent. The supernatant was added into three sterile falcons. Petroleum ether was used to extract the non-polar compounds with the alternative ratios (petroleum ether: extract = 2:1; 1:1; 1:2). The mixtures were mixed vigorously. After collection of the petroleum ether fractions, the remaining extract was alternatively added with chloroform with the suitable ratio of chloroform: supernatant (2:1; 1:1; 1:2). Repeating the same steps as petroleum ether, the mixtures were mixed vigorously. The collected chloroform fractions were evaporated and then the powder was dissolved in distilled water for antimicrobial activity and cytotoxicity. The remaining extract continued to be purified with methanol and applied the same process. The fraction was used for antimicrobial activity test.

#### 3. Results and discussion

**3.1.** Result:

#### a. Phylogeny tree analysis

Using Blast search, isolated fungi had 28 SrRNA showing 99% homology to *Talaromyces purpurogenus* IAM15392. *T. purpurogenus* was classified formerly in *Penicillaceae* because of their close root. However, *T. purpurogenus* was classified in different category nowadays. The isolated fungi lie within the tree of *T. purpurogenus*. The isolated fungi was identified as *Talaromyces purpurogenus* AD1 (Figure 1).



Figure 1: Phylogeny tree of isolated microoganism using neighbour-joining

### b. Antimicrobial activities

*T. purpurogenus* expressed antimicrobial activity to all strains of tested pathogens. Results of antimicrobial test were recorded in table 1.

Table 1: Diameter of inhibition zones resulted from antimicrobial test of *T.purpurogenus* extract with different kinds of pathogens.

	Sal	Pseudo	Serrat	Listeria	Vibrio
	mon	monas	ia	monocyt	parahaem
	ella	aerugi	marce	ogenes	olyticus
	typh	nosa	scens	(17)	(VSA)
	i	( <b>P3</b> )	( <b>P6</b> )		
	( <b>P</b> 2)				
24h	8.07	7.23±0.	6.87±	8.43±0.	5.23±0.25
	±0.1	25	0.32	15	
	5				
<b>48h</b>	9.33	8.77±0.	8.50±	9.60±0.	7.13±0.32
	±0.3	25	0.26	26	
	1				
72h	10.7	12.17±	10.23	12.07±0	11.00±0.26
	3±0.	0.21	±0.15	.49	
	31				
96h	12.2	13.30±	11.10	13.23±0	12.03±0.35
	3±0.	0.36	±0.36	.32	
	21				
120h	13.9	13.43±	12.67	13.97±0	12.97±0.25
	3±0.	0.15	±0.15	.06	
	31				
240h	16.7	16.17±	15.50	16.37±0	15.43±0.21
	7±0.	0.21	±0.21	.15	
	21				
	1				

In *Salmonella typhi*, the weakest antimicrobial activity was expressed after culturing *T. purpurogenus* in 24h that the inhibition zone was smallest and measured about  $8.07\pm0.15$  mm (Table 1). The activity increased until culturing in 240h, showing the highest activity. The inhibition zone was measured about  $16.77\pm0.21$  mm (Table 1).

In *Pseudomonas aeruginosa*, the weakest antimicrobial activity was expressed after culturing *T. purpurogenus* in 24h that the inhibition zone was smallest and measured about  $7.23\pm0.25$  mm (Table 1). The activity increased

© 2018 IJMSIR, All Rights Reserved

until culturing in 240h, showing the highest activity. The inhibition zone was measured about  $16.17\pm0.21$  mm (Table 1).

The weakest antimicrobial activity against *Listeria monocytogenes* was expressed after culturing *T. purpurogenus* in 24h that the inhibition zone was smallest and measured about  $8.43\pm0.15$  mm (Table 2-1). The activity increased until culturing in 240h, showing the highest activity. The inhibition zone was measured about  $16.37\pm0.15$  mm (Table 1).

In *Serratia marcescens*, *T.purpurogenus* showed the weakest antimicrobial activity after 24h of culturing that the inhibition zone was smallest and measured about  $6.87\pm0.32$  mm (Table 2-1). The activity increased until culturing in 240h, expressing the highest activity. The inhibition zone was largest and measured about  $15.50\pm0.21$ mm (Table 1).

*T. purpurogenus* expressed the weakest antimicrobial activity against *Vibrio parahaemolyticus* after 24h of culturing where the inhibition zone measured about  $5.23\pm0.25$  mm (Table 2.1). After 240h of culturing, the activity increased, showing the highest activity where the inhibition zone was measured about  $15.47\pm0.50$  mm (Table 1).

From the results of antimicrobial activities determined in different incubation time that was summarized in Table 1, the analysis of the effects of incubation time on the antimicrobial activities of *T. purpurogenus* was done (Figure 1).





From Figure 2, the results exhibited that antimicrobial activity of the T.purpurogenus continuously increased from 24 hours to 240 hours of culturing. This species showed the best antimicrobial activity at 240 hours with all types of pathogens.

### c. Preliminary purification

Because T. purpurogenus extract showed biological activity, the study tried to perform the preliminary purification for understanding about characteristics of the biological agents. Petroleum ether and chloroform fractions were collected, evaporated and dispersed into distilled water. The powders obtained in petroleunm ether and chloroform fractions could be soluble in water. As a result, petroleum ether and chloroform fractions could contain certain compounds which were soluble in both polar and non-polar solvents.

### d. Antimicrobial test of the purified substance

In order to test which fraction contained the antimicrobial agent, the antimicrobial tests were done.

The partial purified substances in chloroform fraction had the antimicrobial activity with Salmonella typhi (P2) and Listeria monocytogenes (17) (Figure 3). However, it showed no zone of inhibition with Pseudomonas aeruginosa (P3), Serratia marcescens (P6), Vibrio parahaemolyticus (VSA).





Salmonella typhi and Listeria monocytogenes with different purified ratio using chloroform for extraction. From figure 3, it was seen that the inhibition zones which exhibited the antimicrobial activity of the purified substances became larger when the ratio chloroform: supernatant increase from (1:2) to (2:1). In contrast, the inhibition zones of the supernatant after purification become smaller when the ratio: chloroform: supernatant increase with the same ratio.

### e. Cytotoxicity activity on Hela cells

The initial concentration of the stock solution was 1g/mL. It was diluted to the concentration of 100mg/mL in order to perform the test. The test showed no cytotoxic activity on Hela, in contrast, showed high cytotoxic percentage at the concentration of 100 mg/mL (97.29  $\pm$  0.99 %). The extract was diluted into different concentrations to determine the half maximal inhibitory concentration (IC50) on Hela cancer cell line. Result of this test was recorded in Table 2.

Table 2: Cytotoxic activity and the half maximal inhibitory concentration (IC50) of T. purpurogenus on Hela cancer cell lines

Concentration	Cytotoxic percentage (%)				
(mg/mL)	1 <sup>st</sup> time	2 <sup>nd</sup> time	3 <sup>rd</sup> time	Mean ±	24
				•••••	

				SD
100	96.15	97.80	97.92	97.29 $\pm$
				0.99
75	94.00	94.40	93.75	94.05 ±
				0.33
50	58.89	62.80	65.28	$62.32 \pm$
				3.22
25	17.49	17.72	16.50	17.24 ±
				0.65
12.5	3.54	5.97	1.16	3.55 ±
				2.41
IC50	49.02	47.50	48.36	48.29 ±
( <i>mg/mL</i> )				0.76

*T. purpurogenus* showed the weakest cytotoxicity to MCF-7 cancer cell line  $(0.45 \pm 4.30 \%)$  while the strongest cytotoxic percentage was expressed at Hela  $(97.29 \pm 0.99\%)$ . IC50 of *T. purpurogenus* on Hela cancer cell lines was  $48.29 \pm 0.76$  (mg/mL).

#### 3.2. Discussion

YM medium was chosen for culturing *T. purpurogenus* because it is rich in nutrients. YM provided fungus with two important nutrient sources, carbon source (in dextrose, beef extract) and protein source (in beef extract and yeast). In addition, the pH of medium is 6.5, nearly equal to the pH required for culture fungi medium. From figure 1 and table 1, it could be concluded that *T. purpurogenus* was able to produce substances with antimicrobial activity when cultured in YM medium.

As showing in figure 1, more than one antimicrobial agents in *T. purpurogenus* supernatant because the activities did not form a straight curve when culturing from 24h to 240h. In the case of *Pseudomonas*, the activity increased from 24h to 72h, then the activity looked stable until 120h, but sudden increased until 240h. It was meant that there was a complexity of antimicrobial activities in *T. purpurogenus*. To identify the

antimicrobial compounds, more studies should be done so far. During preliminary purification, after using petroleum ether extraction, the antimicrobial activities was not detectable. The supernatant layer stayed below the petroleum ether layer and above the chloroform layer. From figure 3, the results indicated that there are more than one substance with having antimicrobial activity are produced during the time of culturing T.purpurogenus. In fact, the supernatant remained the antimicrobial activities against all pathogens even though the chloroform gave the antimicrobial activities against Salmonlla typhi and Listeria monocytogenes. The more amount of chloroform, the more substances were purified. The inhibition zones were also greater. The inhibition zone of the supernatant became smaller each time the amount of chloroform solvent increase. It means the supernatant has lost substance during the purification using chloroform and that substance has laid in the chloroform layer.

With the concentration 100 mg/mL, the supernatant exhibited high cytotoxic ability with Hela cancer cell line (97.29  $\pm$  0.99 %). IC50 was detected at 48.29  $\pm$  0.76 (mg/mL). The mechanism of new agent will identified so far.

#### 4. Conclusion

YM medium was suitable for culturing *T.purpurogenus* to produce antimicrobial compounds. *T.purpurogenus* will continue to be cultured with effects from other factors like Temperature, pH, Inoculation along with Time to optimize cultural condition for obtaining the highest yield of biological compounds.

Substances laid in chloroform layer had antimicrobial activity. Components as well as molecular structure of these substances will be detected using Thin layer chromatography, High performance liquid chromatography, Infrared spectroscopy and Nuclear magnetic resonance in future research.

*T.purpurogenus* culture extract exhibited cytotoxic effect on Hela cancer cell line. Substances with cytotoxic ability will be purified and detected for molecular formula and structure for further study.

#### References

- Zhao J, Shan T, Mou Y, Zhou L, "Plant-derived bioactive compounds produced by endophytic fungi", Mini Rev Med Chem, vol. 11, no. 2, 2011, p. 159-68.
- Wilson BJ, Wilson CH, "Extraction and preliminary characterization of a hepatotoxic substance from Penicillium rubrum", Journal of Bacteriology, vol. 84, 1962, p. 283–290.
- Moss MO, Robinson FV, Wood AB, Paisley HM, Feeney J, "Rubratoxin B, a proposed structure for a bis-anhydride from Penicillium rubrum Stoll", Nature, 220, 1968, p. 767–770
- Natori S, Sakaki S, Kurata H, Udagawa S, Ichinoe M, Saito M, et al, "Production of rubratoxin B by Penicillium purpurogenum Stoll", Applied Microbiology, 19, 1970, p. 613–617.
- Burnside JE, Sippel WL, Forgacs J, Carll WT, Atwood MB, Doll ER, "A disease of swine and cattle caused by eating moldy corn: Experimental production with pure cultures of molds", American Journal of Veterinary Research, 18, 1957, p. 817–824.
- Lockard VG, Watson SA, Siraj MY, Hayes AW, O'Neal RM, "Rubratoxin B hepatotoxicity: An electron microscopic study", Experimental and Molecular Pathology, 34, 1981, p. 94–109.
- Surjono T, Syrief T, Sudarwati S, Okada K, "Sensitive period for rubratoxin B-induced malformations in ICR mice", Congenital Anomalies, 25, 1985, p. 297–304.
- Engelhardt JA, Carlton WW, Rebar AH, "Rubratoxin B mycotoxicosis in the Syrian hamster", Food and Chemical Toxicology, 25, 1987, p. 685–695.

- Kihara T, Surjono TW, Sakamoto M, Matsuo T, Yasuda Y, Tanimura T, "Effects of prenatal rubratoxin-B exposure on behaviors of mouse offspring", Toxicological Sciences, 61, 2001, p. 368– 373.
- Richer L, Sigalet D, Kneteman N, Shapiro J, Jones A, Scott RB, et al, "Fulminant hepatic failure following ingestion of moldy homemade rhubarb wine", Gastroenterology, 112, 1997, A1366.
- Wang T, Zhang Y, Wang Y, Pei Y, "Anti-tumor effects of Rubratoxin B on cell toxicity, inhibition of cell proliferation, cytotoxic activity and matrix metalloproteinase-2,9", Toxicology in Vitro, 21, 2007, p. 646–650.
- Wada S, Usami I, Umezawa Y, Inoue H, Ohba S, Someno T, et al. "Rubratoxin A specifically and potently inhibits protein phosphatase 2A and suppresses cancer metastasis", Cancer Science, 101(3), 2010, p. 743–50.
- 13. Cunha BA, "Pseudomonas aeruginosa: resistance and therapy", Semin Respir Infect, 2002, p. 231-9.
- 14. Heltberg O, "Nosocomial epidemic of Serratia marcescens septicemia ascribed to contaminated blood transfusion bags", Transfusion, 1997, p. 221-7.
- Lister PD, "Chromosomally-encoded resistance mechanisms of Pseudomonas aeruginosa: therapeutic implications", Am J Pharmacogenomics, 2002, p. 235-43.
- 16. Mills SD, "Comparison of Salmonella typhi and Salmonella typhimurium invasion, intracellular growth and localization in cultured human epithelial cells", Microb Pathog, 1994, p. 409-23.
- Meca G, "Isolation and purification of enniatins A, A(1), B, B(1), produced by Fusarium tricinctum in solid culture, and cytotoxicity effects on Caco-2 cells", Toxicon, 2010, p. 418-24.

© 2018 IJMSIR, All Rights Reserved

- Skehan P, Storeng R, Scudiero D, "New colorimetric cytotoxicity assay for anticancer-drug screening", J Natl Cancer Inst, 1990, p. 1107-12.
- Batra A, Sen A, "Valuation of antimicrobial activity of different solvent extracts of medicinal plant: Melia Azedarach L", International Journal of Current Pharmaceutical Research, 4(2), 2012, p, 67-73.
- 20. S.K.M.Basha RS, "Antimicrobial activity of Indigofera Barberi, a high valued endemic herb of Tirumala Hills", Indian Journal of Fundamental and Applied Life Sciences, 2013, p. 683-687.
- Vichai V, Kirtikara K, "Sulforhodamine B colorimetric assay for cytotoxicity screening", Nat Protoc, 2006, p. 1112-6.