

A Study of Group a Streptococcal Pharyngitis among 3–15-Year-Old Children Attending Clinics for an Acute Sore Throat

Sadhana Joshi¹, Gaurav Parashar²

¹Senior Demonstrator Department of Microbiology, ²Senior Resident Department of ENT

^{1,2} RVRS Medical College, Bhilwara

Corresponding Author: Gaurav Parashar, Senior Resident Department of ENT, RVRS Medical College, Bhilwara

Type of Publication: Original Research Paper

Conflicts of Interest: Nil

Abstract

Background: Group A beta-hemolytic *Streptococcus* (GABHS) is the most important cause of sore throat in children for obvious reasons.

Methods: This was a cross-sectional hospital-based study in the ENT OPD. The study population included all children between ages 3 and 15 years presenting at the OPD.

Results: GABHS was isolated in 70.00% of the children. *Streptococcus viridans* was found in 4 (13.3%) while the remaining 5 (16.67%) were sterile. The age group with the highest proportion of GABHS was age 3–6 years. The youngest age group (3–6 years) also had the highest frequency of *S. viridans* which is a normal flora.

Conclusion: The current study has shown that the proportion of GABHS throat infection is high.

Keywords: Children, Group A beta-hemolytic *Streptococcus*, rheumatic heart disease, sore throat.

Introduction

Group A beta-hemolytic *Streptococcus* (GABHS) is the most important cause of sore throat in children for obvious reasons. It is a causative organism of pharyngitis that is linked to the etiopathogenesis of acute rheumatic

fever (ARF) and rheumatic heart disease (RHD). Post-World War II era saw a lot of efforts, research, and interventions in primary and secondary prevention of RHD leading to a drastic reduction in the incidence of the disease, especially in developed countries. Nevertheless, RHD remains among the most common acquired heart diseases in many countries, especially the developing ones. A previous report from Abeokuta showed that RHD persists as an important cause of heart disease.¹⁻²

Concerning clinical manifestations, children with streptococcal pharyngitis present with fever, sore throat, and nonspecific symptoms such as headache, abdominal pain, and vomiting. The most suggestive physical findings are diffuse redness of the tonsils and tonsillar pillars and soft palate petechial mottling, with or without lymphadenitis and follicular exudates. The benefit of antibiotic treatment of GAS pharyngitis is not limited to accelerating clinical recovery. When initiated within 9 days of illness onset, antibiotic is highly effective for the prevention of ARF. Therefore, it is recommended that antibiotic therapy should not be delayed in children with symptomatic GAS pharyngitis. The drug of choice is penicillin. However, allergy to penicillin precludes its use

in some patients. Alternatives include amoxicillin, erythromycin, azithromycin, clarithromycin, or first-generation cephalosporins. It is important to remember that there could be local variations in antibiotic susceptibility, and this must be put into careful consideration in choosing antibiotics.³⁻⁴

Material and methods

This was a cross-sectional hospital-based study in the ENT OPD. The study population included all children between ages 3 and 15 years presenting at the OPD.

Inclusion and exclusion criteria- Consecutive children aged 3–15 years who presented with sore throat or drooling of saliva and any one of these following signs and symptoms were considered eligible: fever >37.5C, cervical lymphadenopathy, inflamed tonsils, and exudative tonsils. Exclusion criteria were denial of consent and those with significant history of vomiting.

Specimen collection-Throat swabs were collected from all patients recruited into the study using aseptic technique to prevent contamination. A wooden tongue depressor is used to hold the tongue in place. Without touching the sides of the oral cavity or the tongue, a sterile swab stick is used to swab the posterior pharynx and tonsillar arches. The specimens were taken to the laboratory by a research assistant immediately after collection.

Results

Table 1. Socio-demographic variable

Socio-demographic variable		No of cases	Percentage
Gender	Male	17	56.67
	Female	13	43.33
Age groups(Yrs)	3-6	14	46.67
	7-10	11	36.67
	>10	5	16.67

Out of 30 children, 17 (56.67%) were males while 13 (43.33%) were females giving male-to-female ratio of 1.1.31. The mean age of the children was 7.342 ± 3.84 years (range 3–14 years).

GABHS was isolated in 70.00% of the children. *Streptococcus viridans* was found in 4 (13.3%) while the remaining 5 (16.67%) were sterile. The age group with the highest proportion of GABHS was age 3–6 years. The youngest age group (3–6 years) also had the highest frequency of *S. viridans* which is a normal flora.

Discussion

There appears to be a consensus that antibiotic treatment of *S. pyogenes* is an effective primary prevention strategy for ARF and RHD, both of which are its nonsuppurative complications. However, following the drastic reduction of RHD burden, especially in developed countries of the world, there seem to be divided opinions on whether acute sore throat should be investigated and treated with antibiotics as a means of primary prevention of ARF/RHD or not.⁵⁻⁶ Whereas ARF and RHD are rarely seen in the developed countries and therefore weaken the basis for antibiotic treatment, RHD remains a major cause of acquired heart diseases in developing countries and may be nonuniformly distributed even within the same country. As noted earlier in this article, our hospital records, though unpublished, show persistent new cases of ARF/RHD, and therefore, the need to continue to investigate and treat GABHS pharyngitis is key to control of RHD in developing countries.

Sadoh *et al.* in Benin,⁵ Nigeria, reported that 48.72% of the bacterial isolates from throat swabs were beta-hemolytic streptococci, but no Lancefield grouping was done (Group A, B, C, and G are beta-hemolytic).

Another study in Jos, Nigeria, by Mawak *et al.*⁶ reported a much lower figure of 10.45% for *S. pyogenes* among under-five age group.

Conclusion

The current study has shown that the proportion of GABHS throat infection is high.

References

1. Floyd WD. History of hemolytic streptococci. In: Stevens D, Kaplan E, editors. Streptococcal Infections: Clinical Aspects, Microbiology, and Molecular Pathogenesis. New York: Oxford University Press; 2000. p. 1-15.
2. Pruthvish S. Epidemiology of Group A streptococcal infections, rheumatic fever and rheumatic heart diseases and role of WHO. In: Vijayalakshmi IB, editor. Acute Rheumatic Fever and Chronic Rheumatic Heart Diseases. New Delhi: Jaypee Brothers Medical; 2011. p. 35-46
3. Mendis S, Puska P, Norrving B, editors. Global Atlas on Cardiovascular Disease Prevention and Control. Geneva: World Health Organization (in collaboration with the World Heart Federation and World Stroke Organization); 2011.
4. Wilson SE, Chinyere UC, Queennette D. Childhood acquired heart disease in Nigeria: An echocardiographic study from three centres. Afr Health Sci 2014;14:609-16.
5. Sadoh WE, Sadoh AE, Oladipo AO, Okunola OO. Bacterial isolates of tonsillitis and pharyngitis in paediatric casualty setting. J Med Biomed Res 2008;7:37-44.
6. Mawak JD, Ewelike IC, Lar PM, Zumbes HJ. Bacterial etiologic agents associated with upper respiratory tract infections in children (under five

years) attending selected clinics in Jos, Nigeria. Highland Med Res J 2006;4:22-30.