



International Journal of Medical Science and Innovative Research (IJMSIR)

IJMSIR : A Medical Publication Hub Available Online at: www.ijmsir.com

Volume - 2, Issue - 3, May - June - 2017, Page No. : 81 - 83

A Study on Correlation of Degree of Midline Shift on CT scan And Glasgow COMA Scale In Patients of Acute Traumatic Head Injury

Preetam¹, Deepika Meena², Deepak Meena³, Manish Kumar Meena⁴, G L Meena⁵

^{1,5}Department of Radiodiagnosis, SP Medical College & Associate Group of PBM Hospitals, Bikaner.

² Rajasthan Dental College Jaipur, Rajasthan, India

³Mahtama Gandhi Dental College Jaipur, Rajasthan, India

⁴ S.N. Medical College, Jodhpur, Rajasthan, India

Correspondence Author: Dr G.L.Meena, Department of Radiodiagnosis, SP Medical College & Associate Group of PBM Hospitals, Bikaner, Rajasthan India

Conflicts of interest: None to Declare

Abstract

Background: Traumatic head injury affects up to 2% of population per year, constitute the major cause of death.

Methods: A study was conducted on patients with acute traumatic head injury. Most common and important complication of traumatic head injury is the development of an increased intracranial pressure resulting in midline shift. The larger the amount of the midline shift on CT scan the poorer will be the outcome of traumatic head injury.

Results: External injury of scalpel is seen in 90% of cases, blackening of eye in 40% & vomiting in 50 %. Cerebral contusion (40%) was the most common CT scan finding followed by depressed fracture (30%), subdural hematoma (20%) than extradural hematoma (10%). Hemorrhagic contusion was the most common CT scan finding irrespective of GCS score.

Conclusions: The increased degree of midline shift in patients with head injuries by CT scan was related to the severity of head injury (GCS= 3-12) and was significantly related to poor final clinical outcome.

Keywords: CT scan, Glasgow coma scale, Acute traumatic head injury.

Introduction

Traumatic head injury affects up to 2% of population per year, constitute the major cause of death.

Most common and important complication of traumatic head injury is the development of an increased intracranial pressure resulting in midline shift.

The larger the amount of the midline shift on CT scan the poorer will be the outcome of traumatic head injury. Other variables such as Glasgow coma scale have been subsequently introduced to build more complex and accurate prognostic model. In Glasgow coma scale it was found that confident prediction could be made only after 24 hours^{1,2}.

Methods

This study was conducted in department of Radiology SPMC Bikaner during April 2016 to March 2017.

The criteria for inclusion of patients in the present study were

1) All traumatic head injury patients.

2) All patients who had initial cranial Computed Tomography (CT) scan after head injury.

In all the patients CT scan was reviewed separately from clinical information. The degree of midline shifting was divided into three categories; no shifting, midline shifting up to 10 mm & midline shifting greater than 10 mm.

Corresponding Author: Dr. G. L. Meena, ijmsir, Volume-2 Issue-3 Page No. 81 - 83

age 8

Dr. G. L. Meena, et al. International Journal of Medical Sciences and Innovative Research (IJMSIR)

Clinical information such as age, gender, mechanism of head injury, Glasgow coma scale & clinical outcome were collected.

Severity of head injury is classified into 3 subgroups; mild degree (GCS=15), moderate degree (GCS=13-14), severe degree (GCS=3-12).

Results

Table 1: Age	wise	distribution	in	case of	TBL
--------------	------	--------------	----	---------	-----

Age (years)	No. of cases
20-29	6
30-39	7
40-49	8
50-59	9
60-69	12
70-79	4
80-89	4
Total	50

Most of patients(24%) 60-69 years agegroup.

Sex	No. of cases
Male	35
Female	15
Total	100
Total	100

Out of 100 cases included in the study 70 % are males while females constituted only 30% showing male preponderance in acute TBI.

Table 3: Distribution of etiologies in acute TBI.

Etiology	No. of cases
RTA	40
Fall	11
Assault	9
Total	50

80% cases were RTA.

External injury of scalpel is seen in 90% of cases, blackening of eye in 40% & vomiting in 50 %. Cerebral contusion (40%) was the most common CT scan finding followed by depressed fracture (30%), subdural hematoma (20%) than extradural hematoma (10%). Hemorrhagic contusion was the most common CT scan finding irrespective of GCS score. In patients with GCS 3-5 other outcome findings are extradural hematoma, subdural hematoma, & depressed fracture. In patients with GCS 6-8 other common findings are extradural hematoma, depressed fracture & hemorrhagic contusion. In patients with GCS 9-12 other common findings were hemorrhagic contusion, depressed fracture & intra cerebral hematoma. In patients with GCS 13-15 other common findings were depressed fracture, hemorrhagic contusion.

Discussion

RTA is the most common etiological factor in TBI cases. Third and fourth decade of life is the commonest age group with male gender because they are involved in economic and social life³⁻⁶. It is found that children and young adults are most commonly involved in trauma ³⁻⁷. In our study there is increase in the incidence of TBI from pediatric age group to young adults followed by a plateau which constitutes the age group of 40- 70 years which is followed by decline in incidence constituting the age group of above 70 years.

The present study found that the presence of midline shift especially with SDH was significant. This meant that the outcome would be poorest if the midline shift with SDH compared to other lesion in patients with brain injury.

Conclusion

The increased degree of midline shift in patients with head injuries by CT scan was related to the severity of head injury (GCS= 3-12) and was significantly related to poor final clinical outcome.

References

- Azian AA, Nurulazman AA, Shuaib L, Mahayidin M, Ariff AR, Naing NN, et al. Computed tomography of the brain in predicting outcome of traumatic intracranial haemorrhage in Malaysian patients. Acta Neurochir (Wien). 2001;143(7):711- 20.
- Adams JH. The neuropathology of head injuries. In: Vinken PJ, Bruyn GW, eds. Handbook of Clinical Neurology. Vol. 23. New York: Elsevier; 1975: 35-65.
- Kelly AB, Zimmerman RD, Snow RB, et al: Head trauma: Comparison of MR and CT— experience in 100 patients. AJNR 9:699-708, 1988.
- Sklar EM, Quencer RM, Bowen BC, et al: Magnetic resonance applications in cerebral injury. Radiol Clin North Am 30:353-366, 1992.
- Yealy DM, Hogan DE: Imaging after head trauma. Who needs what? Emerg Med Clin N Am 9:707-717, 1991.
- Borezuk P: Predictors of intracranial injury in patients with mild head trauma. Ann Emerg Med 25:731-736, 1995.
- Reinus WR, Wippold FJ, Erickson KK: Practical selection criteria for noncontrast cranial computed tomography in patients with head trauma. Ann Emerg Med 22:1148-1155, 1993