

**A Study on the Lipid Profile of Hypertensive Patients**

<sup>1</sup>Dr. Kali Charan Sonkriwal, Senior Specialist (Medicine), Govt. District Hospital, Kekri, Ajmer

<sup>2</sup>Dr. Sudhir Mathur, Principal Specialist (Medicine), Govt. District Hospital, Kekri, Ajmer

**Corresponding Author:** Dr. Sudhir Mathur, Principal Specialist (Medicine), Govt. District Hospital, Kekri, Ajmer

**Citation this Article:** Dr. Kali Charan Sonkriwal, Dr. Sudhir Mathur, “A Study on the Lipid Profile of Hypertensive Patients”, IJMSIR- August - 2020, Vol – 5, Issue - 4, P. No. 136 – 139.

**Type of Publication:** Original Research Article

**Conflicts of Interest:** Nil

**Abstract**

**Background:** To study the lipid profile in hypertensive patients

**Methods:** The present study was carried out on a total of 100 hypertensive patients attending our Hospital and 100 age and sex matched healthy controls. Twelve hour fasting lipid analysis was done for Serum triglycerides (TG), total cholesterol (TC), High Density Lipoprotein (HDL), Low Density Lipoprotein (LDL).

**Results:** The Mean serum total cholesterol values were highly significant in hypertensive subjects ( $236.12 \pm 34.26$  mg/dL) as compared to the healthy control subjects ( $156.23 \pm 12.10$  mg/dL). The mean serum TG level was  $106.23 \pm 16.12$  mg/dL in healthy control subjects, and  $214.12 \pm 34.21$  mg/dL in hypertensive patients. This difference was highly significant. The mean values for HDL was  $28.41 \pm 6.03$  mg/dL for hypertensive subjects whereas  $41.23 \pm 4.12$  mg/dl for healthy control subjects, respectively. The Mean serum LDL values were highly significant in hypertensive subjects ( $162.32 \pm 31.24$  mg/dL) as compared to the healthy control subjects ( $94.21 \pm 17.21$  mg/dL)

**Conclusion:** All the lipid profile like cholesterol, LDL, TG and HDL were derange in the hypertensive than those in the healthy controls.

**Keywords:** Hypertension, Lipid profile, Disease.

**Introduction**

Arterial hypertension is a major cause of morbidity and mortality because of its association with coronary heart disease, cerebrovascular disease and renal disease. The extent of target organ involvement (i.e. heart, brain and kidneys) determines outcome. North American studies have shown that hypertension is a major contributor to 500 000 strokes (250 000 deaths) and 1 000 000 myocardial infarctions (500 000 deaths) per annum. National surveys continue to reveal that hypertension is often not detected and, where diagnosed, is often inadequately treated. Among hypertensive patients, only 25% appear to be well controlled. This is particularly true of isolated systolic hypertension. Yet the prevalence of isolated systolic hypertension increases with age. Indeed, the proportion of subjects suffering from isolated systolic hypertension, as opposed to systolic and diastolic hypertension, increases from 20% in the under 40 yr to 80% in the 60–69 yr old, and to 95% in those >80 yr.

There is increasing emphasis on the risk associated with systolic hypertension as the level of systolic pressure is a good predictor of coronary and cerebrovascular risk, especially in the elderly. Treatment of systolic hypertension with its wide pulse pressure is effective in terms of control of blood pressure and reduced morbidity, especially in older patients with high risk profile.<sup>1,2</sup>

The blood pressure however, is not the only determinant of cardiovascular damage and the propensity of hypertensive patients to develop target organ damage is markedly influenced by coexisting risk factors such as age, sex, smoking, obesity, dyslipidemia and others.<sup>3-4</sup>

Dyslipidaemia and hypertension are the commonest risk factors for coronary artery disease (CAD). Recent reports show that borderline hypertension (systolic BP 130-139 and/or diastolic BP 85-89 mmHg) and Stage I hypertension carry a significant cardiovascular risk and there is a need to reduce this blood pressure<sup>5</sup> The reported prevalence of hypertension varies around the world, with the lowest prevalence in rural India (3.4% in men and 6.8% in women) and highest in Poland (68.9% in men and 72.5% in women).<sup>6</sup> It has been found that men have a higher prevalence of hypertension than women although this changes later in life with substantial increase in the number of females with hypertension after the age of 50 years. Dyslipidaemia (hyperlipidaemia), which is associated with hypertension, has been recognised as independent risk factor for cardiovascular disease, a leading diagnosis for visits to physicians and cause of death.<sup>7,8</sup>

### Materials and Methods

Patients who diagnosed as cases of essential hypertension based on history and on Antihypertensive

Medication were included. The selected subjects were further grouped as:

Group 1: Healthy control subjects (n=100). It was ensured by routine examination that all the subjects were healthy and there were no signs and symptoms of hypertensive and other disease.

Group 2: Hypertensive subjects (n=100). It included the clinically established patients of hypertension. There blood pressure is in range of systolic blood pressure (>140 mmHg) and diastolic blood pressure (>90mmHg) and have no symptoms of diabetes mellitus.

An informed consent was taken from all the healthy control subjects and patients, under study apprising them the nature and objective of the study. All subjects were studied as outpatient. Participant's examination included interviews for medical and nutritional history. After on overnight fast of 10-12 hours, fasting blood samples were collected. Blood samples were drawn from antecubital vein of each subject by using aseptic technique. The blood was collected in plain tubes for lipid parameters respectively. Serum was separated after centrifugation and analysed.

Lipid profile measured following methods

- Serum total cholesterol: was measured by Enzymatic method Normal serum cholesterol: 150-250 mg/dl
- Serum HDL cholesterol: was measured by "Phosphotungstate method. Normal HDL – Cholesterol: 30 – 70 mg/dl.
- Serum LDL cholesterol: If the value of Triglycerides is known, LDL-cholesterol can be calculated based on Friedewald's equation.
- Serum Triglycerides: was measured by enzymatic colorimetric method Normal Serum Triglycerides: Male: 60-165 mg/dl Female: 40-140 mg/dl.

**Results**

Table 1: Socio-demographic variable

Socio-demographic variable	GROUP-1	GROUP-2	P-value
Age	48.26±6.21	49.12±6.45	>0.05
M:F	68:32	70:30	>0.05

Socio-demographic variable between both groups were comparable.

Table 2: Blood pressure

BP	GROUP-1	GROUP-2	P-value
SBP	116.23±5.14	148.24±4.08	<0.05
DBP	78.24±4.16	103.20±3.16	<0.05

The mean systolic blood pressure of the healthy controls and hypertensive subjects in the present study was 116.23±5.14 mmHg and 148.24±4.08 mmHg . The mean diastolic blood pressure of the healthy controls and hypertensive subjects in the present study was 78.24±4.16 mmHg and 103.20±3.16mmHg respectively.

Table 3: Lipid profile

Lipid profile	Group -1	Group -2	P-value
TC (mg/dl)	156.23±12.10	236.12±34.26	<0.001
LDL (mg/dl)	94.21±17.21	162.32±31.24	<0.001
HDL (mg/dl)	41.23±4.12	28.41±6.03	<0.001
TG (mg/dl)	106.23±16.12	214.12±34.21	<0.001

The Mean serum total cholesterol values were highly significant in hypertensive subjects (236.12±34.26 mg/dL) as compared to the healthy control subjects (156.23±12.10 mg/dL). The mean serum TG level was

106.23±16.12 mg/dL in healthy control subjects, and 214.12±34.21 mg/dL in hypertensive patients. This difference was highly significant. The mean values for HDL was 28.41±6.03 mg/dL for hypertensive subjects whereas 41.23±4.12 mg/dl for healthy control subjects, respectively. The Mean serum LDL values were highly significant in hypertensive subjects (162.32±31.24 mg/dL) as compared to the healthy control subjects (94.21±17.21mg/dL).

**Discussion**

The Mean serum total cholesterol values were highly significant in hypertensive subjects (236.12±34.26 mg/dL) as compared to the healthy control subjects (156.23±12.10 mg/dL). The mean serum TG level was 106.23±16.12 mg/dL in healthy control subjects, and 214.12±34.21 mg/dL in hypertensive patients. This difference was highly significant. The mean values for HDL was 28.41±6.03 mg/dL for hypertensive subjects whereas 41.23±4.12 mg/dl for healthy control subjects, respectively. The Mean serum LDL values were highly significant in hypertensive subjects (162.32±31.24 mg/dL) as compared to the healthy control subjects (94.21±17.21mg/dL)The significantly higher plasma total cholesterol, triglycerides and LDL-cholesterol in the hypertensive than in the normotensive patients in the present study is in corroboration with earlier studies.<sup>9-12</sup>

In accordance to our study, Saha MS et al (2006) also reported a statistically highly significant relation in serum TG level in hypertensive subjects (184.77±5.97 mg/dL) as compared to the healthy controls (142.73±6.68 mg/dL).<sup>13</sup>

Kumar NL et al (2010) reported a statistically highly significant relation in serum TG level in hypertensive subjects (180.88±68.5 mg/dL) as compared to the healthy controls (114.7±17.62 mg/dL).<sup>14</sup>

## Conclusion

All the lipid profile like cholesterol, LDL, TG and HDL were derange in the hypertensive than those in the healthy controls.

## References

1. WHO. World Health Organisation- International Society of Hypertension Guideline for the management of hypertension (Guideline Sub-Committee). *J Hypertens* 1999; 151-183.
2. The sixth Reports of the Joint National Committee on Detection, Evaluation and Treatment of High Blood Pressure (JNC VI). *Arch Intern Med* 1997; 157: 2413-2446.
3. WHO. Classification of hypertension. Report of WHO Scientific Group, Technical Report Series 1978; 657: 87-95.
4. Gupta R: trends in hypertension epidemiology in india. *J Hum Hypertens*. 2004; 18:73-78.
5. Kearney P M, Whelton M, Reynolds K, Whelton Pk, H J, Worldwide prevalence of hypertension: a systematic review. *J Hypertens*. 2004 Jan; 22(1):11-9.
6. Gordon H. hypertensive vascular disease. In: Eugene Braunwald et. al. (ed). *Harrison's Principles of Internal Medicine* 15th edition. McGraw-New York 2000; Pp 141-1430.
7. Kadiri S. Current concepts in the management of hypertension. *Dokita* 1999; 26: 93-96.
8. Shahid A, Zuberi SJ, Hasnain N. Lipid pattern in healthy subjects. *Pak J Med Res* 1985; 24: 33-7.
9. Youmbissi TJ, Djoumessi S, Nouedoui C. Profile lipidique d'un group d'hypertendus camerounais noir Africains. *Medicine d'Afrique Noire* 2001; 31: 114-118.
10. Ahaneku JE, Nwosu MC, Ahaneku GI, Okugba PC. Utilisation of Clinical chemistry tests with special reference to lipid profile in disease management in a Nigeria setting. *East Afr Med J* 1999; 76:172-175.
11. Mgonda YM, Ramaiya KL, Swai ABM, Mc-Larty DG, George KM, Alberti M. Insulin resistance and hypertension in non-obese Africans in Tanzania. *Hypertension* 1998; 31: 114-118.
12. Jarikre AE, Dim DC, Ajuluchukwu JNA. Plasma lipid levels in Nigerian hypertensives: the gender factor. *Nig Qtr J Hosp Med* 1996; 6: 293-298.
13. Saha MS, Sana NK and Ranajit Kumar Shaha, Serum lipid profile of hypertensive patients in the northern region of the Bangladesh..*J. Bio-Sci*; 2006;14: 93-98.
14. Kumar NL, Deepthi J, Rao YN, Deedi MK, Study of lipid profile, serum magnesium and blood glucose in hypertension. *Biology and Medicine*; 2010; 2(1): 6-16.