

**A Study To Assess Predictive Role of C-Reactive Protein In Early Pregnancy among Women attending Obstetrics and Gynaecology OPD in SMS Medical College, Jaipur**

Dr. Purva Sharma<sup>1</sup>, Dr. Mohan Lal Meena<sup>2</sup>

<sup>1</sup>Resident Doctor, <sup>2</sup>Senior Professor

Department of Obstetrics & Gynaecology, SMS Medical College & Attached Group of Hospitals, Jaipur, Rajasthan

**Corresponding Author:** Dr. Mohan Lal Meena, Senior Professor, Department of Obstetrics & Gynaecology, SMS Medical College & Attached Group of Hospitals, Jaipur, Rajasthan.

**Type of Publication:** Original Research Article

**Conflicts Of Interest:** Nil

**Abstract**

**Background:** High concentrations of maternal C-reactive protein have been associated with adverse pregnancy outcome, and premature uterine contraction may be predicted by elevated levels of C-reactive protein. This may ultimately be simple and cost-effective enough to introduce as a low-risk screening program.

**Methods:** Hospital based comparative analysis was conducted at Department of Obstetrics and Gynaecology, SMS Medical College and Associated Hospitals, Jaipur. Women with early pregnancy upto 14 weeks with either abdominal pain or vaginal bleeding or suspected extrauterine pregnancy. C-reactive protein (CRP) quantitative estimation is done by turbidimetric method. Collected samples were sent to a designated lab of our hospital.

**Results:** The mean c-reactive protein level in cases was 2.24 mg/dl with min-max value ranging from 0.84-3.92mg/dl while in controls mean c-reactive protein value came to be 9.04mg/dl with min-max range from 3.28-24.8mg/dl. The difference between the two groups is significant as p value is less than 0.001.

**Conclusion** Our results of significantly increased CRP levels in normal pregnancy and a clear association between CRP and normal pregnancy, support the clinical application of this diagnostic tool in early pregnancy, especially as a predictor of abnormal first trimester pregnancies.

**Keywords:** CRP, Pregnancy, Women.

**Introduction**

Early pregnancy is described as first 12 weeks from beginning of the last menstrual period to the end of first three months. A pregnancy may end in a live birth, abortion or an ectopic in rare cases.<sup>1,2</sup>

In first trimester, raised CRP levels have been reported and more recently it was shown that women with higher CRP levels at 9-13 weeks are more likely to develop gestational diabetes mellitus and pre-eclampsia.<sup>3</sup>

It is not possible to assess the maternal fetal interface in ongoing or threatened pregnancies directly. Hence, the interest in circulating factors such as cytokines.<sup>4</sup>

One of the most important markers of an inflammatory response is C- reactive protein (CRP), first discovered in 1930 as an acute phase protein that reacted with the c-polysaccharide of pneumococcus bacteria. CRP does

not cross the placental barrier and therefore, will be useful in diagnosing infections in newborns.<sup>5</sup>

Recently, it has been shown that CRP is present in amniotic fluid and fetal urine, and the elevated levels are associated with adverse pregnancy outcome. These results demonstrate that the human placenta produces and releases CRP, like other placental proteins, mainly into the maternal circulation.

### Material And Method

**Type of Study:** Hospital based comparative analysis.

**Study Design:** Prospective Study

**Place of Study:** Department of Obstetrics and Gynaecology, SMS Medical College and Associated Hospitals, Jaipur.

**Duration:** From June 2018 to July 2019 when desired sample size was achieved (after taking the approval from Institutional Review Board and Ethical committee).

**Study Participants:** The study included women with early pregnancy upto 14 weeks divided further as:

**Case:** Women with early pregnancy upto 14 weeks with either abdominal pain or vaginal bleeding or suspected extrauterine pregnancy.

**Control:** First trimester pregnancy upto 14 weeks

**Sampling Procedure:** All eligible consecutive cases were enrolled in cases and control group till desired sample size was attained respectively.

### Inclusion Criteria

Patient who gave written and informed consent.

Women with known first trimester pregnancy (upto 14 weeks of pregnancy) who were referred to our department with or without abdominal pain, vaginal bleeding and suspected extrauterine pregnancies.

### Exclusion Criteria

- Pregnant women who had already been treated with methotrexate for ectopic pregnancy

- Women with a known chronic or acute inflammatory condition (e.g. inflammatory bowel disease, or arthritis, lupus vasculitis, heart diseases, pneumonia, burns, trauma).

- Steroid/NSAID users or taking drugs like thiazolidinone and statins.

### Methodology

- All eligible patients fulfilling inclusion criteria were explained about nature and purpose of the study.

- After taking their informed and written consent, detail history, general and systemic examination were done.

- Patients venous blood samples were collected for CRP in plain vial along with routine blood investigations and USG for foetal well being.

- C-reactive protein (CRP) quantitative estimation is done by turbidimetric method. Collected samples were sent to a designated lab of our hospital.

- All information and reports were recorded on a predesigned proforma and were entered in Microsoft Excel sheet to prepare master chart.

### Observations and Results

Table No. 1: Distribution of Study Participants According to Age

Age Group	Case		Control	
	No.	%	No.	%
20-22 yrs	13	21.66	16	26.67
23-25 yrs	38	63.33	32	53.33
26-27 yrs.	9	15.00	12	20.00
Total	60	100.00	60	100.00
Mean±SD	23.75±1.64		23.82±1.83	
P value	0.833 (NS)			

According to table number 1 mean age of study participants was 23.75 yrs which was found similar to that in control group. Hence, there is no statistical

difference (p =0.8) among the groups as per age difference.

Table No. 2: Distribution of study participants according to gestation age

Weeks of Gestation	Case		Control		Total	
	No.	No.	%	%	No.	%
5-10	51	85.00	51	85.00	102	85.00
11-14	9	15.00	9	15.00	18	15.00
Total	60	100.00	60	60.00	120.00	100.00

Chi-square = 0.065 with 1 degree of freedom; P = 0.798 (NS)

According to table 9 it was observed that 85% of cases and 85% of controls were from 5-10 weeks of gestation; and 15% cases and controls had 11-14wks of gestation. Therefore two groups were similar by period of gestation and p value was 0.7 (NS).

Table No. 3: Comparison of mean Serum C-Reactive Protein level between Case and Control

CRP Level (mg/dl)	Case	Control
Sample size	60	60
Mean ± SD	2.24±0.75	9.04±5.01
Median	2	7
Min-Max	0.84-3.92	3.28-24.8
P value	P<0.001	

The mean c-reactive protein level in cases 2.24 with min-max value ranging from 0.84-3.92mg/dl while in controls mean c-reactive protein value came to be 9.04 with min-max range from 3.28-24.8mg/dl.

Hence mean c-reactive protein levels were higher among controls showing that a higher c-reactive protein level was seen among pregnant women in early gestation with normal intrauterine pregnancy when compared to abnormal presentations in first trimester of pregnancy.

## Discussion

As per table no. 10 the mean c-reactive protein level in cases 2.24 with min-max value ranging from 0.84-3.92 mg/dl while in controls mean c-reactive protein value came to be 9.04 with min-max range from 3.28-24.8mg/dl.

Hence mean c-reactive protein levels were higher among controls showing that a higher c-reactive protein level was seen among pregnant women in early gestation with normal intrauterine pregnancy when compared to abnormal ones. Our study results are in line with similar studies conducted by : Watts et al 1991<sup>6</sup> reported 7-9 mg/l CRP levels for pregnant women not in labour starting from 22 weeks of gestation compared to non-pregnant population.

GP Sacks et al 2004<sup>7</sup> observed that pregnant women had significantly higher CRP levels (median 3.68 mg/l) than those who were not pregnant (median 1.495 mg/l, P < 0.0001), a difference that persisted after excluding potential confounding variables. Boggess KA, Lieff (2005)<sup>8</sup> study was to determine the relationship between maternal inflammation and first or second trimester pregnancy loss compared maternal serum C-reactive protein concentration between women with a pregnancy loss at < 21 weeks gestation to control women without gestational diabetes or preeclampsia who delivered at term. Median serum C-reactive protein concentration was significantly higher in controls compared with all cases (3.2 versus 0.5 microg/mL; p <0.001). Also studies conducted by Pitiphat et al (2005)<sup>9</sup> and Lohsoonthorn et al (2007)<sup>10</sup> found there is statistically significant association between CRP concentrations in early pregnancy >8mg/l and subsequent preterm delivery, with odds ratios of 2.55 and 2.04; respectively in two studies

## Conclusion

Our results of significantly increased CRP levels in normal pregnancy and a clear association between CRP and normal pregnancy, support the clinical application of this diagnostic tool in early pregnancy, especially as a predictor of abnormal first trimester pregnancies.

## References

1. Williams Obstetrics: 25<sup>th</sup> edition chapter 18 abortion pg 348-349; chapter 5 implantation and placenta pg 95,97
2. National Institute for Health and Clinical Excellence. Ectopic pregnancy and miscarriage: diagnosis and initial management in early pregnancy of ectopic pregnancy and miscarriage. NICE Clinical Guideline 154. Manchester (UK): NICE; 2012. Available at: <http://www.nice.org.uk/guidance/cg154/resources/guidance-ectopic-pregnancy-and-miscarriage-pdf>. Retrieved January 20, 2015.
3. Wilcox AJ, Weinberg CR, O'Connor JF, Baird DD, Schlatterer JP, Canfield RE, et al. Incidence of early loss of pregnancy. *N Engl J Med* 1988;319:189-94.
4. Black S, Kushner I, Samols D. C-reactive protein. *J Biol Chem* (2004) 279(47):48487-90.10.1074/jbc.R400025200
5. Gewurz H, Mold C, Siegel J, Fiedel B. C-reactive protein and the acute phase response. *Adv Intern Med* (1982) 27:345-72.
6. C Reactive protein in Normal Pregnancy Watts DH, Krohn 199
7. G.P. Sacks. Maternal C-reactive protein levels are raised at 4 weeks gestation. *Human Reproduction*, 2004; 19( 4): pp-1025-1030, 2004.
8. Boggess KA, Lieff S, Murtha AP, Moss K, Jared H, Beck J, et al. 2005. Maternal serum C-reactive protein concentration early in pregnancy and subsequent pregnancy loss
9. Pitiphat W, Gillman MW, Joshipura KJ, et al. Plasma C-Reactive Protein in Early Pregnancy and Preterm Delivery. *Am J Epidemiol*. 2005;162:1108-1113.
10. LohsoonthornVQC, Williams MA. Maternal serum C-reactiveProtein concentrations in early pregnancy and subsequent risk of preterm delivery. *ClinBiochem*. 2007 Mar;40(5-6):330-335